Mon. Not. R. Astron. Soc. **000**, 000–000 (0000)   Printed 5 February 2008   (MN LaTeX style file v2.2)

Kinetically Dominated FRII Radio Sources

Brian Punsly
4014 Emerald Street No.116, Torrance CA, USA 90503 and
International Center for Relativistic Astrophysics, I.C.R.A.,University of Rome La Sapienza, I-00185 Roma, Italy
E-mail: brian.m.punsly@L-3com.com or brian.punsly@gte.net

arXiv:astro-ph/0610042v1 2 Oct 2006

5 February 2008

ABSTRACT

The existence of FR II objects that are kinetically dominated, the jet kinetic luminosity, Q, is larger than the total thermal luminosity (IR to X-ray) of the accretion flow, L_{bol}, is of profound theoretical interest. Such objects are not expected in most theoretical models of the central engine of radio loud AGN. Thus, establishing such a class of objects is an important diagnostic for filtering through the myriad of theoretical possibilities. This paper attempts to establish a class of quasars that have existed in a state of kinetic dominance, $R(t) \equiv Q(t)/L_{bol}(t) > 1$, at some epoch, t. It is argued that the 10 quasars in this article with a long term time average $Q(t)$, $\overline{Q}$, that exceed L_{Edd} are likely to have satisfied the condition $R(t) > 1$ either presently or in the past based on the rarity of $L_{bol} > L_{Edd}$ quasars. Finally, the existence of these sources is discussed in the context of the theory of the central engine.

Key words: quasars: general — galaxies: jets — galaxies: active — accretion disks — black holes.

1 INTRODUCTION

The connection between accretion flow parameters and radio jet power is mysterious. Most quasars are radio quiet, however $\approx 10\%$ of optically selected quasars are radio loud and more importantly about 2% of quasars have powerful FRII radio lobes deVries et al. (2006). The lobe emission is the signature that the time averaged jet kinetic luminosity, $\overline{Q}$, is enormous, $\overline{Q} > 10^{44}$ergs/sec Punsly (2001). It is not known how powerful the quasar jet can be relative to the thermal luminosity from accretion, L_{bol}. Understanding the limits of jet power can help reveal the physical nature of the quasar central engine. The primary obstacle in this exercise is that in order for L_{bol} and Q to be estimated contemporaneously necessitates that $Q(t)$ be derived from parsec scale radio jet observations and such efforts are plagued by poorly estimated Doppler enhancement factors (raised to the fourth power) and the results are often grossly inaccurate Punsly (2005); Punsly and Tingay (2006). For example, the estimates of blazar jet power in Celotti et al (1997)); Wang et al. (2004) indicate that $R(t) \equiv Q(t)/L_{bol}(t) > 1$ AGN are fairly common, but the results are skewed by poorly constrained Doppler factors Punsly and Tingay (2005). Alternatively, the time averaged jet power $\overline{Q}$ can be estimated far more accurately from the isotropic properties of the extended emission. Some $\overline{R} \equiv \overline{Q}/L_{bol} > 1$ sources were found in Punsly and Tingay (2006). Unfortunately the $\overline{Q}$ estimate is not contemporaneous with the L_{bol} data, so one can not say if the sources presently satisfy or ever satisfied $R(t) > 1$. Grossly inaccurate measurements of Q are really of no value, so we must concentrate on the $\overline{Q}$ estimates and combine this with other information in order to establish a class of $R(t) > 1$ sources. In section 3, a subsample of 10 FR II AGN with $\overline{Q}_{Edd} \equiv \overline{Q}/L_{Edd} > 1$, is argued to be comprised of $R(t) > 1$ sources.

2 ESTIMATION TECHNIQUES

In this section, we review the standard estimation techniques used in the following analysis. The more information that is known about the large scale radio structure such as the radio spectral index across the lobe and high resolution X-ray contours, the more sophisticated and presumably more accurate an estimate that can be obtained for the energy flux delivered to the lobes from the jet Punsly (2005); Birzan et al (2004); Punsly (2001). Unfortunately, such detailed information does not exist for most radio sources and we need an expedience that is helpful for studying large samples. Such a method that allows one to convert 151 MHz flux densities, F_{151} (measured in Jy), into estimates of $\overline{Q}$ (measured in ergs/s), was developed in Willott et al. (1999); Blundell and Rawlings (2000), the result is captured by the formula derived in Punsly (2005):

$$\overline{Q} \approx 1.1 \times 10^{45} \left[X^{1+\alpha} Z^2 F_{151}\right]^{\frac{6}{7}} \text{ergs/sec}\,, \qquad (1)$$

$$Z \equiv 3.31 - (3.65) \times$$

$$\left[X^4 - 0.203X^3 + 0.749X^2 + 0.444X + 0.205\right]^{-0.125} \quad (2)$$

where $X \equiv 1+z$, F_{151} is the total optically thin flux density from the lobes (i.e., **contributions from Doppler boosted jets or radio cores are removed**). In this paper we adopt the following cosmological parameters: H_0=70 km/s/Mpc, $\Omega_\Lambda = 0.7$ and $\Omega_m = 0.3$. We define the radio spectral index, α, as $F_\nu \propto \nu^{-\alpha}$. The formula is most accurate for large classical double radio sources, thus we do not consider sources with a linear size of less than 20 kpc. Alternatively, one can also use the independently derived isotropic estimator from Punsly (2005)

$$\overline{Q} \approx 5.7 \times 10^{44} (1+z)^{1+\alpha} Z^2 F_{151}\,\text{ergs/sec}$$
$$\alpha \approx 1\ . \quad (3)$$

Even though one can use the jet emission from the parsec scale radio core to estimate, Q more contemporaneously with the accretion flow emission as in Celotti et al (1997)), such estimates are prone to be very inaccurate. One is observing a very small amount of "waste energy" such as X-ray or optical emission as the powerful radio jet propagates away from the source. One must then try to figure out the fraction of Q that is converted to waste energy in this region. Typically, the X-ray, radio and optical regions are observed with different spatial resolution, so it is unclear if one is detecting the same physical region on parsec scales as one synthesizes the broad band data. In cases in which there is sufficient broad band flux to make an estimate, one is plagued with the further ambiguity of determining the Doppler factor of the relativistic jet. This is a critical obstacle because the luminosity from an unresolved region scales with the Doppler factor to the fourth power Lind and Blandford (1985). More shortcomings of this method are discussed in Punsly and Tingay (2005) and a published example in which the Q is apparently miscalculated using radio core properties by three orders of magnitude is discussed explicitly. Thus, the most accurate estimates of Q should use an isotropic estimator such as the radio lobe flux in (1), even though it is just a time average.

The total bolometric luminosity of the accretion flow, L_{bol}, is the thermal emission from the accretion flow (IR to X-ray), including any radiation in broad emission lines. To estimate, L_{bol}, we use the composite SED in table 2 of Punsly and Tingay (2006), which see for details. This spectrum, in combination with the broad emission lines, represents the "typical" radiative signature of a strong accretion flow onto a black hole. This signature is empirical and it is independent of all theoretical models of the accretion flow. If $L(\nu)_{\rm obs}$ is the observed spectral luminosity at the quasar rest frame frequency, ν, then L_{bol} is estimated as

$$L_{bol} = 1.35 \frac{\nu L(\nu)_{\rm obs}}{\nu L(\nu)_{\rm com}} \times 10^{46}\text{ergs/sec}\ , \quad (4)$$

where $L(\nu)_{\rm com}$ is the spectral luminosity from the composite SED.

Finally, we list the equations used to estimate the central black hole mass, M_{bh}. The reader should consult the references for details. We estimate M_{bh} from the line widths, F(H β) or F(C IV), Vestergaard and Peterson (2006) or F(Mg II), Kong et al. (2006), where F() means "FWHM of"

$$M_{bh}(H\beta) =$$
$$10^{6.91\pm0.02}\left[\left(\frac{F(H\beta)}{1000\text{km/s}}\right)^2 \left(\frac{\lambda L_\lambda(5100\mathring{A})}{10^{44}\text{ergs/s}}\right)^{0.50}\right]\ , \quad (5)$$

$$M_{bh}(\text{CIV}) =$$
$$10^{6.66\pm0.01}\left[\left(\frac{F(\text{CIV})}{1000\text{km/s}}\right)^2 \left(\frac{\lambda L_\lambda(1350\mathring{A})}{10^{44}\text{ergs/s}}\right)^{0.53}\right]\ , \quad (6)$$

$$M_{bh}(\text{MgII}) =$$
$$10^{6.53}\left[\left(\frac{F(\text{MgII})}{1000\text{km/s}}\right)^2 \left(\frac{\lambda L_\lambda(3000\mathring{A})}{10^{44}\text{ergs/s}}\right)^{0.58\pm0.10}\right]\ . \quad (7)$$

3 KINETICALLY DOMINATED SOURCES

Table 1 is comprised of $\overline{Q}_{Edd} \equiv \overline{Q}/L_{Edd} > 1$ QSOs that were found by cross-correlating all existing VLA, MERLIN and ATCA radio maps with the absolute visual magnitudes and broad line widths of the QSOs in Veron-Cetty and Veron (1991). Sources that were discovered by this method with $\overline{R} > 4$ were previously reported in Punsly and Tingay (2006), which see for more details on the sample selection. A more in depth analysis showed that 3C 216 and 3C 455 also belong in the table. However, as mentioned previously, due to the lack of simultaneity between jet ejection and UV emission, the Punsly and Tingay (2006) sample does not establish the $R(t) > 1$ condition. In subsection (3.1) it is argued that the $\overline{Q}/L_{Edd} > 1$ jets in table 1 satisfy $R(t) > 1$, for some t. The first column is the source name followed by the redshift. Column 3 has two estimates of $\overline{Q}$ from (1) on the left and (3) on the right, separated by a slash. Column 4 is L_{bol} estimated from (4) at the rest frame frequency in column 6. Column 5 is $\overline{R}$. The last three columns are L_{bol}/L_{Edd}, $\overline{Q}_{Edd}$ and the references. If more than one observed frequency was available, a second estimate for L_{bol} was provided for that source so that one can assess the error in using a composite SED to approximate L_{bol}.

The QSO, 3C 455, requires some elaboration. This object has a narrow Hβ (FWHM = 620 km/s), yet the nuclear luminosity is far stronger than a narrow line radio galaxy and its magnitude is more typical of a strong Seyfert 1 galaxy, hence its historical classification as a quasar Gelderman and Whittle (1994). Implicit in the first estimate in table 1 is that the object is a narrow line Seyfert 1, but Hβ appears weak compared to [OIII]λ5007 (Hβ/[OIII]=0.2 and Hβ/[OIII]> 0.33 in narrow line Seyfert1's) because the narrow line region is greatly enhanced by emission line gas that is excited by the jet and is aligned with the radio structure deVries et al. (1999). In support of this interpretation, there is an elongated patch of diffuse emission enveloping the kpc radio jet axis between the two lowest contours of the HST image in Lehnert et al. (1999) with a flux approximately equal to the narrow line flux in the F702W filter Gelderman and Whittle (1994); deVries et al. (1999). Alternatively, one can interpret the object as a narrow line radio galaxy and M_{bh} can be estimated from the galactic host bulge luminosity. There are a variety of fits to the M_{bh} - host bulge luminosity relation, but the fits of McClure and Dunlop (2002) offer the smallest scatter since they are based on large samples. The relation that is most relevant is the best fit estimator derived from 72 AGN in

Table 1. FR II Quasars with Super Eddington Jets

Source	z	$\overline{Q}$ 10^{45}ergs/s	L_{bol} 10^{45}ergs/s	$\overline{R}$	freq (10^{15} Hz)	L_{bol}/L_{Edd}	$\overline{Q}_{Edd}$	ref
3C 216	0.670	15.1/14.1	≈ 0.12	≈ 120	0.71/1.16	$0.05-0.1$	$3.3-10$	1
3C 455	0.543	7.13/5.04	0.38	18.7/13.3	0.94	1.42	26.7/18.9	2
		7.13/5.04	0.38	18.7/13.3	0.94	0.07	1.33/0.94	3
3C 82	2.878	155.4/183.8	14.5	10.7/12.7	0.014	0.106	1.14/1.35	4
		155.4/183.8	25.0	6.22/7.35	1.67	0.245	1.52/1.80	4
3C 9	2.009	148.3/174	25.0	5.93/6.96	1.67	0.264	1.57/1.85	5
		148.3/174	38.8	3.82/4.49	0.0078	0.324	1.24/1.46	6
4C 25.21	2.686	59.3/59.7	11.6	5.11/5.15	1.14	0.198	1.02/1.02	5
PKS 1018-42	1.28	63.9/65.2	19.3	3.31/3.38	1.37	0.428	1.42/1.45	7
		63.9/65.2	14.7	4.35/4.45	1.37	0.326	1.42/1.45	7
4C 04.81	2.594	103.8/148	35.8	2.90/4.13	2.30	0.459	1.33/1.90	5
3C 196	0.871	73.5/87.0	31.6	2.33/2.76	1.53	3.04	7.10/8.41	8
		73.5/87.0	31.6	2.33/2.76	1.53	0.238	0.66/0.56	9
3C 14	1.469	52.38/51.68	32.6	1.61/1.59	1.00	0.604	1.05/1.03	10
3C 270.1	1.519	65.1/66.6	48.2	1.35/1.38	2.07	0.844	1.14/1.17	5

1. see Punsly (2007), 2. continuum and FWHM from Gelderman and Whittle (1994), M_{bh} from eqn (5), 3. M_{bh} from bulge luminosity estimate in eqn (8), 4. L_{bol} and FWHM raw data from Semenov et al. (2004), M_{bh} from eqn (6), 5. L_{bol} and FWHM from Barthel et al. (1990), M_{bh} from eqn (6), 6. continuum from Meisenheimer et al. (2001), FWHM from Barthel et al. (1990),M_{bh} from eqn(6), 7. Punsly and Tingay (2006), M_{bh} from eqn(7), 8. continuum and FWHM from Lawrence et al. (1996), M_{bh} from eqn (5), 9. continuum and FWHM from Lawrence et al. (1996), M_{bh} from eqn (7), 10. continuum and FWHM from Aars et al (2005), M_{bh} from eqn (7)

table 3 of McClure and Dunlop (2002),

$$\log(M_{bh}/M_{\odot}) = (-0.46 \pm 0.03)M_R - (2.55 \pm 0.72)\ .(8)$$

Thus, we need to find M_R for the galactic bulge in order to utilize (8). The HST image is taken with the F702W filter and the apparent magnitude of the galactic bulge (after subtraction of the nuclear core and extended narrow line flux) is $m_{702W} = 21.53$ deVries et al. (1999). The m_{702W} magnitudes are approximately the Cousins R magnitudes. The transformation to the Cousins R magnitude along with the k-correction that is given in Fukujita et al. (1995) yields $M_R = -21.5$ and from (8), $M_{bh} = 4.2 \times 10^7 M_{\odot}$. This is the basis for the second estimate in table 1.

3.1 Super Eddington Jets

A value of $\overline{Q}_{Edd} > 1$ suggests that even if the jet central engine is currently in a low state then at some epoch during the lifetime of the source it must have had $R(t) > 1$. For example, if PKS 1018-42 is not now nor has ever been kinetically dominated then the on average $\overline{L}_{bol} > 1.45 L_{Edd}$ for the lifetime of the source. However, since $\overline{Q} = (1/T)\int_0^T Q(t)dt$, the peak values of the instantaneous $Q(t)$, $\max_t [Q(t)] > 1.45 L_{Edd}$. Thus, one expects that $\max_t [L_{bol}(t)]$ would have to exceed $1.45 L_{Edd}$ by a significant amount at certain epochs in order for the PKS 1018-42 to have always been in an $R(t) < 1$ state. This is inconsistent with the magnitudes of the peaks in the duty cycle of quasar $L_{bol}(t)/L_{Edd}$ based on our current knowledge, see Boroson (2002); Ho (2005). More importantly, the estimates of 268 AGN in Punsly (2007) utilized the same estimators, eqns. (5)-(7), used here and $L_{bol}/L_{Edd} < 1$ for all the broad line AGN. Furthermore, note that most of the QSOs in table 1 have prodigious accretion rates and $\overline{R} > 1$ is not a consequence of the quasars being in a state of suppressed accretion. The most likely explanation of the data in table 1 is that these sources experienced epochs in which $R(t) \sim 1 - 10$ as opposed to the alternative explanation that the broad line sources had protracted phases in which $L_{bol}(t)/L_{Edd} \sim 1 - 10$.

3.2 The Validity of the Estimates

Since the data set is small, the analysis of the data is particularly sensitive to the integrity of the estimates in table 1. The main drawback to the argument above is that the estimates of M_{bh} could be off by a large amount in an individual object, Vestergaard and Peterson (2006), and these sources appear to have $\overline{Q}_{Edd} > 1$ merely as a result of these errors. If this were the fundamental explanation of the $\overline{Q}_{Edd} > 1$ estimates then a statistical anomaly must also present, the $0.5 \leqslant \overline{Q}_{Edd} \leqslant 1.0$ QSOs just below the threshold of table 1 never have M_{bh} over estimated (i.e. there are no false negatives for the criteria $\overline{Q}_{Edd} > 1$). In subsection 3.2.3, it is shown that this anomalous requirement conflicts with the data used to create the virial mass estimates. Furthermore, there could also be errors of smaller magnitude associated with the estimates of $\overline{Q}$ Punsly (2005).

3.2.1 Suppressing Error Propagation

The possibility of these types of errors in table 1 is dealt with by introducing as many different sets of observational data as possible in order to obtain independent estimates of the same quantities. Thus, any results that are way "out of family" like $\overline{Q}_{Edd} = 26.7$ for 3C 455 can be flagged as unreliable. In order to expose the largest potential source of error, M_{bh}, is computed from as many different spectral bands and emission lines as possible (generally two entries in table 1). Similarly, we compute $\overline{Q}$ from two different estimators. Furthermore, we estimate all potentially reddened QSOs by finding an IR flux estimate to reveal hidden AGN UV luminosity (this will affect (5)-(7)). For example, 3C 190 fails to meet the $\overline{Q}_{Edd} > 1$ criteria because it has considerable hidden AGN UV emission, see table 3 of Punsly and Tingay



Figure 1. A histogram of the distribution of $\log(\overline{Q}_{Edd})$ for the 3CRR QSOs $\geqslant 20$ kpc in linear extent. The center of each bin is indicated on the horizontal axis. All of the sources in the two bin on the far right satisfy $\overline{Q}_{Edd} > 1$. The mean value of $\overline{Q}_{Edd} = 0.52 \pm 0.9$

(2006). The errors might still persist and the potential effects are discussed in the following two subsections.

3.2.2 3CRR Quasars

One way to assess the role of the statistical scatter in the estimators is to study the complete 3CRR sample. We estimated $\overline{Q}_{Edd}$ for every QSO > 20 kpc (remember, (1) and (3) are not accurate for linear sizes less than 20 kpc). Figure 1 is a histogram of the distribution of $\log(\overline{Q}_{Edd})$. Every effort was made to average away errant estimates. Every available broad line FWHM, Lawrence et al. (1996); Brotherton et al (1994); Wills et al. (1995); Aars et al (2005); Gu et al. (2001); Marziani et al (2003); Barthel et al. (1990); Kuraszkiewicz et al (2004), and continuum flux density, Lawrence et al. (1996); Wills et al. (1995); Aars et al (2005); Barthel et al. (1990); Kuraszkiewicz et al (2004); Meisenheimer et al. (2001); Simpson and Rawlings (2000), was gathered for each source from the literature and every combination of these was used to estimate numerous values of M_{bh} from (5)-(7). These values were then averaged for each source. This average M_{bh} and the average $\overline{Q}$ from (1) and (3) were used to compute the "best estimate" of $\overline{Q}_{Edd}$ for each source. These estimates were then assembled in the histogram of figure 1. This histogram shows that the $\overline{Q}_{Edd} \geq 1$ QSOs are just the high end of a smooth distribution of $\overline{Q}_{Edd}$ for the complete sample and are not outliers. Over 30% of the QSOs have $\overline{Q}_{Edd} > 0.5$. This method of assigning $\overline{Q}_{Edd} > 1$ to a QSO is very conservative. A single large estimate for M_{bh} will swamp the other smaller M_{bh} estimates and suppress the large $\overline{Q}_{Edd}$ values (such as in 3C 196, 3C 190 and 3C 455). If one were to average the $\overline{Q}_{Edd}$ values instead then large M_{bh} would be suppressed and there would be four 3CRR QSO with $\overline{Q}_{Edd} > 4$.

3.2.3 An Analysis of the Distribution of Errors

There are two major facts that allow a clear interpretation of the effect of the errors in the estimation techniques on the $\overline{Q}_{Edd} \sim 1$ QSOs in table 1 and figure 1. Firstly, the detailed investigation of 3C 216 in Punsly (2007) illustrates the existence of $\overline{Q}_{Edd} > 1$ QSOs very convincingly. The $\overline{Q}_{Edd} > 3.3$ condition in 3C 216 is verified by three independent estimators of M_{bh}, F(Hβ), F(Mg II), and the host bulge luminosity. Secondly, there is no evidence that the scatter in the best fit estimators is skewed asymmetrically to low values of M_{bh} Vestergaard and Peterson (2006); Kong et al. (2006). Comparisons to reverberation mapping based estimates indicate that the errors are symmetrically distributed about the best fit estimators, eg. figure 9 of Vestergaard and Peterson (2006). If not for these facts, one could take a pessimistic view and plausibly argue that the $\overline{Q}_{Edd} > 1$ sources in table 1 and figure 1 are an artifact of selecting those sources in which the errors associated with the line width estimation techniques have underestimated M_{bh}. This argument implicitly assumes that the high end of the $\log(\overline{Q}_{Edd})$ distribution represents the variance in the line width estimators, not the actual spread in M_{bh}. However, the example of 3C 216 demonstrates that $\overline{Q}_{Edd} > 1$ sources exist, indicating that the more reasonable explanation of figure 1 is that the errors in the estimates are randomly distributed and they are not skewed preferentially to produce low M_{bh}. In particular, some of the sources with $\overline{Q}_{Edd} > 1$ are probably actually, $\overline{Q}_{Edd} < 1$ sources, but a similar number of the $0.5 < \overline{Q}_{Edd} < 1$ sources are actually $\overline{Q}_{Edd} > 1$ sources.

It is worth expanding the discussion of the $\overline{Q}_{Edd} > 1$ sources beyond the most conservative of claims made above. This condition is far stronger than $Q(t)_{\max} > Q_{Edd}$ and in general indicates episodic values of $Q(t)$ considerably in excess of Q_{Edd}. More realistically based on the discussion of section 3.1, all the sources that have $\overline{Q}_{Edd} > 0.5$ almost certainly had episodes in which $Q_{Edd} > 1$ and the $R(t) > 1$ condition was satisfied. Considering that the median value in figure 1 is $\overline{Q}_{Edd} = 0.26$, $\sim 0.3 - 0.5$ of the the 3C sources were likely to have been kinetically dominated at some time in their past. This claim can not be extended to the FR II population as a whole, but is a consequence of the fact that the 3C QSOs typically reside at the high end of the steep extended luminosity distribution for radio loud quasars.

4 DISCUSSION

The article demonstrates that many FR II sources are likely to exist in a state of $R(t) > 1$. The $\overline{Q}_{Edd} > 1$ jets must satisfy $\max_t[R(t)] > 1$, or else the accretion disks of the broad line AGN would episodically be in an un-physical state of $L_{bol}(t)/L_{Edd} \gg 1$. The Mid-IR 3C sample of Ogle et al (2006) indicated that $\sim 1/2$ of the FR II narrow line radio galaxies had no hidden quasar. Based on the discussion above, these are likely candidates to be in a state of $R(t) > 1$.

As a consequence of the near independence of the UV spectrum on the radio state, deVries et al. (2006); Corbin and Francis (1994), it has been argued that the black hole and not the accretion disk is the power source for FR II jets Semenov et al. (2004). A large scale magnetic flux trapped within the central vortex of an accretion disk can produce relativistic jets, as seen in the MHD numerical simulations of Hawley and Krolik (2006); McKinney (2005); McKinney and Gammie (2004). Yet, as discussed in Punsly

(2007) these simulations are restricted to a maximum R value, $R_{max} < 0.5$. The magnetic field line configuration leading to a relativistic jet is created and maintained by "pinning" the poloidal flux to the event horizon by a strong accretion flow in conjunction with an incredibly massive and unobserved enveloping wind that transports $> 1 M_{\odot}$/yr of relativistically hot protonic gas at 0.3c-0.4c for an FR II quasar Punsly (2007). The black hole actually gains energy over time due to the intense accretion flow (and associated energy influx) required to compress a large magnetic flux onto the small surface area of the event horizon. Thus, a jet with large Q requires an intense accretion flow that will radiate profusely for the radiative efficiency predicted in these simulations, Bekwith et al (2006), and $R_{\max} \sim 0.1$

Alternatively, the black hole energy extraction model of Punsly (2001) and references therein that was numerically realized in Semenov et al. (2004) is based on large scale magnetic flux that threads the equatorial plane of the ergosphere (the active region) of a black hole Penrose (1969). The theoretical Q values are 2 orders of magnitude larger than for flux pinned on the event horizon because the surface area of the equatorial plane in the ergosphere is ~ 10 times larger than the surface area of the horizon that is threaded by magnetic flux in simulations of Hawley and Krolik (2006); McKinney (2005); McKinney and Gammie (2004) for rapidly spinning black holes (parameterized by $a/M \approx 1$, where "a" is the angular momentum per unit mass of the black hole in geometrized units), and the jet power scales like the surface area squared Semenov et al. (2004). These solutions attain a value, $R_{\max}$, that was calculated in Punsly (2007, 2001),

$$R_{\max} \approx 11 \frac{0.3}{\epsilon} (a/M)^{10.5} \; , \; 0.90 < a/M < 0.99 \; , \qquad (9)$$

where ϵ is the radiative efficiency of the accretion flow, $L_{bol} \equiv \epsilon (dM/dt)$ and (dM/dt) is the accretion rate from the disk to the black hole. The maximum efficiency from a thin disk is $\epsilon \approx 0.3$ Novikov and Thorne (1973). For high L_{bol} objects, luminous quasars, one expects ϵ to be near maximal and $a/M \approx 1$ (see Bardeen (1970); Elvis et al. (2002)), thus $R_{\max} \sim 10$ which would explain the high Q episodes required to maintain a long term time average of $\overline{Q}_{Edd} > 1$.

These arguments seem to indicate that the essential element that separates a radio quiet QSO from one that launches a strong FR II jet with $R \sim 1$ is the formation of strong vertical magnetic flux in the equatorial plane of the ergosphere. This is prevented in the simulations of Hawley and Krolik (2006); McKinney (2005); McKinney and Gammie (2004) as a very weak seed magnetic field is swept up in a single MHD fluid and compressed onto the black hole. Alternatively, Spruit and Uzdensky (2005) have investigated low angular momentum flux tubes that evolve separately from the bulk of the MHD fluid. In the two-fluid dynamics, the strong flux tubes "swim" relative to the bulk MHD fluid and slowly move into the ergosphere. This could be an essential piece of missing dynamics in the present codes.

REFERENCES

Aars et al 2005, AJ **130** 23

Antonucci, R.J. 1993, Annu. Rev. Astron. Astrophys. **31** 473

Barthel, P., Tytler, D., Thompson, B. 1990, Astron. and Astrophys. Sup. **82** 339

Bardeen, J. 1970 Nature **226** 64

Bekwith, K., Hawley J., Kroilk, J. 2006, astro-ph 0605295

Birzan, L. et al 2004, ApJ **607** 800

Blundell, K., Rawlings, S. 2000, AJ **119** 1111

Boroson, T. 2002, ApJ **565** 78

Brotherton, M. et al 1994, ApJ **423** 131

Celotti, A., Padovani and Ghisellini, G. 1997, MNRAS **286** 415

Corbin, M. and Francis, P. 1994, AJ **108** 2016

deVries, W., Becker, R., White, R. 2006, AJ **131** 666

deVries, W., O'Dea, C., Baum, S. 1999, ApJ **526** 27

Elvis, M., Risaliti, G., Zamorani, G. 2002, ApJL **565** L75

Fukujita, M., Shumasaku, K., Ichikawa, T. 1995, PASP **107** 945

Gelderman, R. and Whittle, M. ApJS **91** 491

Gu, M., Cao, X., Jiang, D.R. 2001, MNRAS **327** 111

Hawley, J., Krolik, K. 2006, ApJ in press astro-ph/0512227

Ho, L. 2005 In: **From X-ray Binaries to Quasars: Black Hole Accretion on All Mass Scales** ed. by Maccarone,T., Fender,R. and Ho, L.C. (Dordrecht: Kluwer) p.219 astro-ph/0504643

Kong, M., Wu, X.-B., Wang, R., Han, J. 2006, Chin.J.Astron.Astrophys. in press astro-ph/0512609

Kuraszkiewicz et al 2004, ApJS **150** 165

Lawrence, C. et al 1996, ApJS **107** 541

Lehnert, M. et al 1999, ApJS **123** 351

Lind, K., Blandford, R. 1985, ApJ **295** 358

Marziani, P. 2003, ApJS **145** 199

McClure, R., Dunlop, J. 2002, MNRAS **331** 795

McKinney, J. and Gammie, C. 2004, ApJ **611** 977

McKinney, J. 2004, ApJL **630**5

Meisenheimer, K. et al 2001 A& A **372**719

I.D. Novikov, K.S. Thorne 1973 In: *Black Holes*, ed. by C. DeWitt and B.S. DeWitt (Gordon and Breach, New York) p. 344

Ogle, P., Whysong, D. and Antonucci, R. 2006, ApJ **647** 161

Penrose, R. 1969, Nuovo Cimento **1** 252

Punsly, B. 2001, *Black Hole Gravitohydromagnetics* (Springer-Verlag, New York)

Punsly, B. 2005, ApJL **623** L12

Punsly, B. 2007, to appear in ApJL, astro-ph/0609204

Punsly, B., Tingay, S. 2005, ApJL **633** L89

Punsly, B., Tingay, S. 2006, ApJL in press

Semenov, V. Dyadechkin, S., Punsly, B. 2004, Science **305** 978

Simpson, C. and Rawlings, S. 2000, MNRAS **317** 1023

Smith, H., Sinrad, H. 1980, ApJ **236** 419

Spruit, H. and Uzdensky, D., ApJ **629** 965

Veron-Cetty, M.P., Veron, M. 1991, *A Catalogue of Quasars and Active Nuclei* 5th edition, European Southern Observatory Scientific Report No. 10

Vestergaard, M. and Peterson, B. 2006, ApJ in press astro-ph/0601303

Wang, J.-M., Luo, B, Ho, L. 2004, ApJL **615** 9

Willott, C., Rawlings, S., Blundell, K., Lacy, M. 1999, MNRAS **309** 1017

Wills, B. et al 1995, ApJ **437** 139